


WHAT IS CLAIMED IS:

1. A compound of Formula (I):




5

(I)

wherein:

L_1 is a group of formula (a):

10



(a)

wherein:

A is an aryl or a heteroaryl ring;

15


B'' is -O-;

R^x is alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, acylamino, aminoacyloxy, aryl, carboxyalkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, heteroaralkyl, alkylsulfonyl, or alkylsulfinyl;


20

R^1 is hydrogen or alkyl;


R^2 is Het, or is selected from a group consisting of formula (i), (ii), and (iii):



(i)



(ii)



(iii)

wherein:

----- is an optional double bond;

n_1 is an integer of from 1 to 4;

25

n_2 is an integer of from 1 to 3;

V is -CH-, -O-, -S(O) n_3 - (where n_3 is an integer of from 0 to 2), or -NR⁴-

(wherein R⁴ is hydrogen, alkyl, substituted alkyl, aryl, or heteroaryl);

“Het” is a heteroaryl ring which optionally attaches (a) to a linker;

R³ is hydrogen, alkyl, halo, amino, substituted amino, -OR^a (where R^a is hydrogen, alkyl, or acyl), or a covalent bond attaching (a) to a linker;

- 5 R⁵ is hydrogen, alkyl, halo, amino, substituted amino, -OR^b (where R^b is hydrogen or alkyl), aryl, aralkyl, heteroaralkyl, or a covalent bond attaching (a) to a linker;

- R⁶, R⁷, and R⁸ are, independently of each other, hydrogen, halo, hydroxy, alkoxy, haloalkoxy, carboxy, alkoxycarbonyl, alkyl optionally substituted with one, 10 two or three substituents selected from halo, hydroxy, carboxy, alkoxycarbonyl, alkylthio, alkylsulfonyl, amino, substituted amino, or a covalent bond attaching (a) to a linker;

K is a bond or an alkylene group;

- K" is a bond, -C(O)-, -S(O)_{n4}- (where n₄ is an integer of from 0 to 2), or an 15 alkylene group optionally substituted with a hydroxyl group; and

B is heterocycloamino or heteroarylamino, which optionally attaches (a) to a linker;


provided that at least one of the R⁵, R⁶, R⁷, R⁸, “Het”, heterocycloamino or heteroarylamino groups attaches (a) to a linker;

- 20 X is a linker; and

L₂ is an organic group comprising at least one primary, secondary or tertiary amine; or a pharmaceutically acceptable salt; or prodrug thereof.

2. The compound of claim 1 wherein L₂ is a group selected from a group 25 consisting of:

- (i) a group of formula (b):



(b)

wherein:

D" is alkylene;

D is $-NR^{31}R^{32}$, $-N^+(R^{33}R^{34}R^{35})$ or $-OR^{32}$ where R^{31} , R^{33} , and R^{34} are,
independently of each other, hydrogen, alkyl, or aralkyl; and R^{32} and R^{35} represent a
5 covalent bond attaching (b) to a linker;


R^{27} is hydrogen, halo, nitro, cyano, hydroxy, alkoxy, carboxy,
alkoxycarbonyl, acyl, thio, alkylthio, alkylsulfonyl, alkylsulfinyl, sulfonamido,
alkylsulfonamido, carbamoyl, thiocarbamoyl, mono or dialkylcarbamoyl, amino,
mono- or dialkylamino, aryl, aryloxy, arylthio, heteroaryl, heteraryloxy,
10 heteroarylthio, heterocyclyl, heterocyclyloxy, aralkyl, heteroaralkyl, or alkyl
optionally substituted with one, two or three substituents selected from halo,
hydroxy, carboxy, alkoxycarbonyl, alkylthio, alkylsulfonyl, amino, or substituted
amino;

R^{28} is hydrogen, halo, nitro, cyano, hydroxy, alkoxy, carboxy,
15 alkoxycarbonyl, acyl, thio, alkylthio, alkylsulfonyl, alkylsulfinyl, sulfonamido,
alkylsulfonamido, carbamoyl, thiocarbamoyl, mono or dialkylcarbamoyl, amino,
mono- or dialkylamino, or alkyl optionally substituted with one, two, or three
substituents selected from halo, hydroxy, carboxy, alkoxycarbonyl, alkylthio,
alkylsulfonyl, amino, or substituted amino;

20 R^{29} and R^{30} are, independently of each other, hydrogen, alkyl, haloalkyl,
halo, nitro, cyano, hydroxy, alkoxy, alkoxycarbonyl, acyl, thio, alkylthio, amino,
mono- or dialkylamino; or

one of R^{27} , R^{28} , R^{29} , or R^{30} together with the adjacent group forms a
methylenedioxy or ethylenedioxy group;

25 (ii) a group of formula (c):



(c)

wherein:

n_{11} is an integer of from 1 to 7;

n_{12} is 0 to 7;

F is $-NR^{40}-$, $-O-$, $-S-$, or $-CHR^{41}-$ (wherein R^{40} and R^{41} are, independently of
5 each other, hydrogen, alkyl, or substituted alkyl);

F'' is a covalent bond, $-OR^{43}$, $-NR^{42}R^{43}$, or $-N^+R^{43}R^{44}R^{45}$ wherein R^{42} is
hydrogen or alkyl, R^{44} and R^{45} are alkyl, and R^{43} is hydrogen, alkyl, or a covalent
bond attaching (c) to a linker;


R^{36} is hydrogen, alkyl, halo, nitro, cyano, hydroxy, alkoxy, carboxy,
10 alkoxy carbonyl, acyl, thio, alkylthio, alkylsulfonyl, alkylsulfinyl, sulfonamido,
alkylsulfonamido, carbamoyl, thiocarbamoyl, mono or dialkylcarbamoyl, amino,
mono- or dialkylamino, aryl, aryloxy, arylthio, heteroaryl, heteraryloxy,
heteroarylthio, heterocyclyl, heterocyclyloxy, aralkyl, heteroaralkyl, or alkyl
optionally substituted with one, two or three substituents selected from halo,
15 hydroxy, carboxy, alkoxy carbonyl, alkylthio, alkylsulfonyl, amino, or substituted
amino;

R^{37} is hydrogen, alkyl, halo, nitro, cyano, hydroxy, alkoxy, alkoxy carbonyl,
acyl, thio, alkylthio, amino, mono- or dialkylamino, aryl, aryloxy, arylthio,
heteroaryl, heteraryloxy, heteroarylthio, heterocyclyl, heterocyclyloxy, aralkyl,
20 heteroaralkyl, or alkyl optionally substituted with one, two or three substituents
selected from halo, hydroxy, carboxy, alkoxy carbonyl, alkylthio, alkylsulfonyl,
amino, or substituted amino; and

R^{38} is hydrogen, alkyl, halo, hydroxy, alkoxy, or a covalent bond attaching
the ligand to a linker provided that at least one of R^{38} and R^{43} attaches (c) to a
25 linker;

R^{39} is hydrogen, alkyl, halo, hydroxy, alkoxy, or substituted alkyl; and

(iii) a group of formula (d) or (e):



wherein:

R⁴⁶ is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, or heterocycle;

R⁴⁷ is alkyl, substituted alkyl, aryl, acyl, heterocycle, or -COOR⁵⁰ where R⁵⁰ is alkyl; or

R⁴⁶ and R⁴⁷ together with the nitrogen atom to which they are attached form heterocycle, which heterocycle, in addition to optionally bearing the optional substituents defined hereinbelow for a heterocycle, can also optionally be substituted with one or more (e.g. 1, 2, 3, or 4) alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, or substituted alkynyl.

R⁴⁸ is a covalent bond that attaches the (d) to a linker; and

R⁴⁹ is alkyl;

or a pharmaceutically acceptable salt; or prodrug thereof.

3. The compound of claim 1 or 2 wherein A is phenyl or pyridyl.

4. The compound of claim 1 or 2 wherein R¹ is hydrogen, methyl, or ethyl.

5. The compound of claim 1 or 2 wherein R² is pyrrolyl, pyridinyl, or imidazolyl.

6. The compound of claim 1 or 2 wherein R² is phenyl.

7. The compound of claim 1 or 2 wherein K is a bond or a methylene group.

8. The compound of claim 1 or 2 wherein K" is a bond.

9. The compound of claim 1 or 2 wherein R^x is alkyl, alkenyl, or alkynyl, each optionally substituted with 1 to 5 alkoxy or fluoro substituents.

10. The compound of claim 1 or 2 wherein R^x is (C₁-C₆)alkyl, (C₂-C₆)alkenyl,

(C₂-C₆)alkynyl, each optionally substituted with 1 to 3 methoxy, ethoxy or fluoro substituents.

11. The compound of claim 1 or 2 wherein R^x is (C₁-C₆)alkyl optionally substituted with 1 to 3 methoxy, ethoxy, or fluoro substituents.

12. The compound of claim 1 wherein R^x is methyl, ethyl, propyl, isopropyl, butyl, isobutyl or secbutyl, optionally substituted with methoxy or ethoxy or with 1 to 3 or fluoro substituents.

10

13. The compound of claim 1 wherein R^x is methyl, ethyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, fluoromethyl, difluoromethyl, trifluoromethyl, trifluoromethoxymethyl, formyl, or acetyl.

14. The compound of claim 1 or 2 wherein R^x is methyl, ethyl, methoxymethyl, fluoromethyl, difluoromethyl, or trifluoromethyl.

15. The compound of claim 1 or 2 wherein B is a heterocycloamino group which attaches (a) to a linker.

20

16. The compound of claim 1 or 2 wherein B is pyrrolidine, piperidine, or hexahydroazepine attaching (a) to a linker.

17. The compound of claim 1 or 2 wherein B is piperidine wherein the nitrogen atom of said piperidine attaches (a) to a linker.

18. The compound of claim 1 or 2 wherein B is piperidin-3-yl or piperidin-4-yl wherein the nitrogen at the 1 position optionally attaches (a) to a linker.


19. The compound of claim 1 wherein B taken together with R^x is 4-methylpiperidine-1,4-diyl.

20. The compound of claim 2 wherein L_2 is a group of formula (d) or (e).


21. The compound of claim 20 wherein: R^{46} is alkyl or substituted alkyl; R^{47} is alkyl, substituted alkyl, or heterocycle; or R^{46} and R^{47} together with the nitrogen atom to which they are attached form heterocycle.

22. The compound of claim 1 or 2 wherein L_2 has any one of the formula A1-A590 shown hereinabove.

23. The compound of claim 1 or 2 wherein L_1 is:




24. The compound of claim 23 wherein L_1 is:



25. The compound of claim 24 wherein the piperidino nitrogen of L_1 is bonded to X.

26. The compound of claim 1 or 2 wherein X is alkylene optionally substituted with one, two, or three hydroxy groups, alkylene wherein one, two or three carbon atoms have been replaced by an oxygen atom, -alkylene-phenylene-alkylene- wherein the phenylene ring is optionally substituted with one or two chloro or fluoro groups.

27. The compound of claim 1 or 2 wherein X is a group of formula:



wherein

5 m is an integer of from 0 to 20;

X^a at each separate occurrence is selected from the group consisting of -O-, -S-, -NR-, -C(O)-, -C(O)O-, -C(O)NR-, -C(S)-, -C(S)O-, -C(S)NR- or a covalent bond where R is as defined below;


10 Z at each separate occurrence is selected from the group consisting of alkylene, substituted alkylene, cycloalkylene, substituted cycloalkylene, alkenylene, substituted alkenylene, alkynylene, substituted alkynylene, cycloalkenylene, substituted cycloalkenylene, arylene, heteroarylene, heterocyclene, or a covalent bond;

15 Y^a and Y^b at each separate occurrence are selected from the group consisting of -O-, -C(O)-, -OC(O)-, -C(O)O-, -NR-, -S(O) $_n$ -, -C(O)NR'-, -NR' C(O)-, -NR' C(O)NR'-, -NR' C(S)NR'-, -C(=NR')-NR'-, -NR'-C(=NR')-, -OC(O)-NR'-, -NR'-C(O)-O-, -N=C(X^a)-NR'-, -NR'-C(X^a)=N-, -P(O)(OR')-O-, -O-P(O)(OR')-, -S(O) $_n$ CR' R''-, -S(O) $_n$ -NR'-, -NR'-S(O) $_n$ -, -S-S-, and a covalent bond; where n is 0, 1 or 2; and R, R' and R'' at each separate occurrence are selected from the group
20 consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, cycloalkenyl, substituted cycloalkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic; provided at least one of X^a , Y^a , Y^b or Z is not a covalent bond.

25 28. The compound of claim 1 or 2 wherein X is an alkylene group having from 3 to 20 carbon atoms; wherein one or more carbon atoms in the alkylene group is optionally replaced with -O-; and wherein the chain is optionally substituted on carbon with one or more hydroxyl.


30 29. The compound of claim 1 or 2 wherein X is nonane-1,9-diyl, octane-1,8-diyl, propane-1,3-diyl, 2-hydroxypropane-1,3-diyl, or 5-oxa-nonane-1,9-diyl.

30. The compound of claim 1 or 2 wherein X has the following formula:




wherein the phenyl ring is optionally substituted with 1, 2, or 3 fluoro groups.

31. The compound of claim 1 or 2 wherein X has one of the following the
5 formula:



32. The compound of claim 2 which is a compound of Formula (Ia):

10




(Ia)

or a pharmaceutically acceptable salt or prodrug thereof.

15

33. The compound of claim 1 or 2 wherein L₂ is a group of formula A234, A363, A364, A153, A28, A324, A329, A562, A87, or A239 as described herein.


34. The compound of claim 1 which is a compound of formula (III):



wherein R², K'', A, K, R¹, X, and L₂ have the values defined in claim 1; or a pharmaceutically acceptable salt or prodrug thereof.

5

35. The compound of claim 34 wherein X is a group of formula:



wherein

10 *m* is an integer of from 0 to 20;

X^a at each separate occurrence is selected from the group consisting of -O-, -S-, -NR-, -C(O)-, -C(O)O-, -C(O)NR-, -C(S)-, -C(S)O-, -C(S)NR- or a covalent bond where R is as defined below;

Z at each separate occurrence is selected from the group consisting of
 15 alkylene, substituted alkylene, cycloalkylene, substituted cycloalkylene, alkenylene, substituted alkenylene, alkynylene, substituted alkynylene, cycloalkenylene, substituted cycloalkenylene, arylene, heteroarylene, heterocyclene, or a covalent bond;


Y^a and Y^b at each separate occurrence are selected from the group consisting
 20 of -O-, -C(O)-, -OC(O)-, -C(O)O-, -NR-, -S(O)_n-, -C(O)NR'-, -NR' C(O)-, -NR' C(O)NR'-, -NR' C(S)NR'-, -C(=NR')-NR'-, -NR'-C(=NR')-, -OC(O)-NR'-, -NR'-C(O)-O-, -N=C(X^a)-NR'-, -NR'-C(X^a)=N-, -P(O)(OR')-O-, -O-P(O)(OR')-, -S(O)_nCR' R''-, -S(O)_n-NR'-, -NR'-S(O)_n-, -S-S-, and a covalent bond; where *n* is
 25 0, 1 or 2; and R, R' and R'' at each separate occurrence are selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, cycloalkenyl, substituted cycloalkenyl, alkynyl,

substituted alkynyl, aryl, heteroaryl and heterocyclic; provided at least one of X^a, Y^a, Y^b or Z is not a covalent bond.

36. The compound of claim 34 wherein X is an alkylene group having from 3 to 20 carbon atoms; wherein one or more carbon atoms in the alkylene group is optionally replaced with -O-; and wherein the chain is optionally substituted on carbon with one or more hydroxyl.


37. The compound of claim 34 wherein X is nonane-1,9-diyl, octane-1,8-diyl, propane-1,3-diyl, 2-hydroxypropane-1,3-diyl, or 5-oxa-nonane-1,9-diyl.

38. The compound of claim 34 wherein X has the following formula:



wherein the phenyl ring is optionally substituted with 1, 2, or 3 fluoro groups.

39. The compound of claim 34 wherein X has one of the following the formula:



40. The compound of claim 2 wherein L_2 is a group of formula (d) wherein R^{46} is a heterocycle, optionally substituted with 1 to 5 substituents independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, and substituted alkynyl; and R^{47} is alkyl, substituted alkyl, acyl, or
5 -COOR⁵⁰.

41. The compound of claim 2 wherein L_2 is a group of formula (d) wherein R^{46} is alkyl that is optionally substituted with from 1 to 5 substituents independently selected from the group consisting of alkoxy, substituted alkoxy,
10 cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, aminoacyl, aminoacyloxy, oxyaminoacyl, cyano, halogen, hydroxyl, keto, thioketo, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, NR^aR^b , wherein R^a and R^b may be
15 the same or different and are chosen from hydrogen, alkyl, substituted alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, and heterocyclic.

42. The compound of claim 2 wherein L_2 is a group of formula (d) wherein R^{46} is 3-piperidinyl, 4-piperidinyl, or 3-pyrrolidinyl, which R^{46} is optionally substituted
20 with 1 to 3 substituents independently selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, cyano, halogen, hydroxyl, keto, thioketo, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy,
25 substituted thioalkoxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, and substituted alkynyl.

43. The compound of claim 2 wherein R^{46} and R^{47} together with the nitrogen
30 atom to which they are attached form a piperidine or pyrrolidine ring which ring is optionally substituted with 1 to 3 substituents independently selected from the

group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, cyano, halogen, hydroxyl, keto, thioketo, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocyclooxy, thiol, thioalkoxy, substituted thioalkoxy, heterocyclic, heterocyclooxy, hydroxyamino, alkoxyamino, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, and substituted alkynyl.

44. The compound of claim 2 wherein R^{46} and R^{47} together with the nitrogen atom to which they are attached form a heterocycle that is an aza-crown ether.

45. The compound of claim 44 wherein the aza-crown ether is 1-aza-12-crown-4, 1-aza-15-crown-5, or 1-aza-18-crown-6.

46. Compound number 1-146 as described in Table A and Table B herein; or a pharmaceutically acceptable salt or prodrug thereof.


47. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of claim 1 or 2.

48. A method of treating a disease mediated by a muscarinic receptor in a mammal, comprising administering to said mammal a therapeutically effective amount of a compound of claim 1 or 2.

49. The method of claim 48 wherein the disease is urinary incontinence chronic pulmonary obstructive disease, asthma, hyper salivation a cognitive disorder, blurred vision, or irritable bowel syndrome.

50. A compound of formula L_1 -H wherein L_1 has the values defined in claim 1; or a salt thereof.

51. The compound of claim 50 which is a compound of formula (V):



(V)

or a salt thereof.

52. A compound of formula R_a-X-L₂ wherein X, and L₂ have the values defined
5 in claim 2; and R_a is a suitable leaving group.

10

15